



Corrigendum to “EASL Clinical Practice Guidelines for the management of patients with decompensated cirrhosis” [J Hepatol 69 (2018) 406–460][☆]

European Association for the Study of the Liver^{**}

It has come to our attention that there is an error in the original manuscript in the section entitled ‘Renal impairment’ on page 429. The sentence ‘Recent studies have suggested that in patients with cirrhosis, in AKI stage 1, SCr <1.5 mg/dl is associated with a worse outcome than an SCr ≥1.5 mg/dl.’ should read

‘Recent studies have suggested that in patients with cirrhosis, in AKI stage 1, SCr ≥1.5 mg/dl is associated with a worse outcome than an SCr <1.5 mg/dl’. We apologise for this mistake and any inconvenience caused.

DOI of original article: <http://dx.doi.org/10.1016/j.jhep.2018.03.024>.

[☆] **Clinical Practice Guideline Panel:** Paolo Angeli (Chair), Mauro Bernardi (Governing Board representative), Panel members: Cándid Villanueva, Claire Francoz, Rajeshwar P. Mookerjee, Jonel Trebicka, Aleksander Krag, Wim Laleman, Pere Gines

^{*} Corresponding author. Address: European Association for the Study of the Liver (EASL), The EASL Building – Home of Hepatology, 7 rue Daubin, CH 1203 Geneva, Switzerland. Tel.: +41 (0) 22 807 03 60; fax: +41 (0) 22 328 07 24.

E-mail address: easloffice@easloffice.eu



ELSEVIER